

EFFECTIVENESS OF EMG BIOFEEDBACK ON IMPROVING HAND FUNCTION IN HEMIPLEGIC STROKE PATIENTS

EFFECTUL BIOFEEDBACKULUI EMG ÎN ÎMBUNĂTĂȚIREA FUNCȚIEI MĂINII LA PACIENȚII HEMIPLEGICI, DUPĂ ACCIDENT VASCULAR CEREBRAL

Maheshwari S Harishchandre¹, Singaravelan R M²

Key words: EMG Biofeedback, Stroke, Hand function, Action Research Arm test, Voluntary control grading scale.

Cuvinte cheie: Biofeedback EMG, AVC, Funcția mâinii, Test al brațului pentru cercetarea acțiunii, Scală de gradare a controlului voluntar.

Abstract.

Objective The purpose of this study was to find out the effect of EMG biofeedback along with conventional physiotherapy on improving hand function in subacute stroke participants.

Design Prospective comparative study.

Setting Department of Neuro Physiotherapy, Pravara Rural Hospital (Tertiary Hospital), Loni, Tal- Rahata, Dist-Ahmednagar, Maharashtra State, India- 413 736.

Methods: This study was conducted between January 2011- November 2011. Thirty participants of both genders, aged between 40-65 and those who met all the eligibility criteria were included. They were divided into two equal groups, Group A Study group received EMG Biofeedback along with conventional physiotherapy and Group B Control group received only conventional physiotherapy. Intervention given over a period of 4 weeks, baseline and post values were assessed by Voluntary Control Grading Scale and Action Research Arm test.

Results: The results of the study revealed that there was a significant difference between both the groups on improved hand voluntary control ($p < 0.01$) and improved hand function ($p < 0.01$) in hemiplegic stroke participants.

Conclusion: The results of this study concluded that EMG Biofeedback along with conventional physiotherapy is effective on improving hand voluntary control and hand function in subacute stage stroke participants over the period of four weeks.

Rezumat.

Obiectiv Scopul acestui studiu a fost aflarea efectului biofeedback-ului EMG împreună cu kinetoterapia convențională asupra îmbunătățirii funcției mâinii la pacienții cu AVC subacut.

Design Studiu comparativ de prospectare

Locație Departamentul de neuro kinetoterapie, Spitalul Rural Pravara (Spital de gradul trei), Loni, Tal-Rahata, Dist-Ahmednagar, Maharashtra, India -413736.

Metode: Studiul s-a desfășurat în perioada ianuarie 2011 – noiembrie 2011. Au fost incluși 30 de pacienți de ambele genuri, cu vârste între 40 și 65 de ani care au corespuns criteriilor. Aceștia au fost împărțiți în două grupuri egale. Grupul de studiu A a beneficiat de Biofeedback EMG și de kinetoterapie convențională în timp ce Grupul de control B a beneficiat numai de kinetoterapie convențională. Intervenția s-a efectuat pe o perioadă de 4 săptămâni, iar evaluările inițială și finală s-au efectuat cu ajutorul Scalei de gradare a controlului voluntar și al Testului brațului pentru cercetarea acțiunii.

Rezultate: Rezultatele studiului au arătat o diferență semnificativă între grupuri în ceea ce privește îmbunătățirea controlului voluntar al mâinii ($p < 0.01$) și îmbunătățirea funcției mâinii ($p < 0.01$) la pacienții hemiplegici după AVC.

Concluzie: Rezultatele studiului au arătat că Biofeedbackul EMG împreună cu kinetoterapia convențională este eficient în îmbunătățirea controlului voluntar al mâinii și a funcției mâinii la pacienții cu AVC în stadiu subacut, pe o perioadă de 4 săptămâni.

¹ Postgraduate Student, College of Physiotherapy, Pravara Institute of Medical Sciences, Loni, Maharashtra State, India - 413 736. Phone: +91-9561745825, +91-9687006553; Fax No: +91-2422-273413; Web: www.pravara.com E-mail: maheshwariharishchandre@gmail.com

² Associate Professor, College of Physiotherapy, Pravara Institute of Medical Sciences, Loni, Maharashtra State, India - 413 736.

Introduction

A stroke, previously known medically as a cerebrovascular accident (CVA), is the rapidly developing loss of brain function(s) due to disturbance in the blood supply to the brain¹. As a result, the affected area of the brain is unable to function, which might result in an inability to move one or more limbs on one side of the body, and various neurological dysfunction like inability to understand or formulate speech.² because of that voluntary control of movement is impaired. According to WHO stroke is the “acute onset of neurological dysfunction due to abnormality in cerebral circulation with resultant signs and symptoms that corresponds to involvement of focal area of brain lasting more than 24 hrs”.³

Stroke is the third leading cause of the death in the western countries and the most common cause of adult disability⁴; of the survivors about 50% will have a significant long term disability⁵. According to W.H.O (16 Nov. 2011) in India incidence of stroke was 130/ 100000 individuals every year⁶. The Middle cerebral artery (MCA) is by far the largest cerebral artery and is the vessel most commonly affected by cerebrovascular accident (CVA). The MCA supplies most of the outer convex brain surface, nearly all the basal ganglia, and the internal capsules⁷. Middle cerebral arterial territory was the most commonly affected (38.9%) followed by posterior cerebral artery 13%, brainstem 13%, anterior cerebral artery in 11.1%, multiple vessel territory 9.3%, small vessel infarcts in 8.4%⁸.

Complete occlusion or blockage, of the middle cerebral artery leads to paralysis of the entire opposite side of the face and body. Partial occlusion can affect only some areas example paralysis of only arm or paralysis of the arm and leg, with sparing the face.^{9,10,11,12} Risk factors for stroke include old age, hypertension (70%), previous stroke or transient ischemic attack (TIA)(10%), diabetes(15% and 49%), high cholesterol(15%), cigarette smoking(9%). High blood pressure is the most important modifiable risk factor of stroke.^{2,13}

Motor deficits are characterized by paralysis (hemiplegia) or weakness (hemiparesis), typically on the side of the body opposite the side of the lesion. Initially flaccid paralysis is present in acute stage and after that in subacute stage muscles goes in spasticity¹⁴. According to Bobath concept there are three stages of stroke. Acute stage, subacute stage and chronic stage. Acute stage is prolonged up to 0-2 week, subacute stage is up to 2 years and chronic stage is more than 2 years.¹⁵

In acute stage wrist muscles are in flaccid state so there is total loss of voluntary movement and immobility which can result in loss of range of motion of wrist flexors. Paralysis of the wrist extensors has a weakening effect upon the flexing ability of the fingers. Paralysis of the finger extensors does not interfere with the interphalangeal joints, so the function is performed by the interossei and lumbricals but if it is in immobile then the contractures of these intrinsic muscles lead to hyperextension of this joints as well as flexion contracture of the metacarpophalangeal joint¹⁶.

Traditionally the techniques which is used for improving hand function includes proprioceptive neuromuscular facilitation technique¹⁷, Task oriented exercises programs, Electrical Muscle Stimulation¹⁸. Recently EMG biofeedback has introduced for improving hand function. As a comprehensive definition, biofeedback is a group of therapeutic procedures that use electronic or electromechanical instruments to properly measure, process & feedback to patients in the form of auditory or abnormal neuromuscular & autonomic activity”¹⁹. EMG biofeedback has been used to improve motor function in patient following stroke. This technique allows participants to alter motor unit activity based on augmented audio & visual feedback information. Training can focus on voluntary inhibition of spastic muscles (e.g. reducing firing frequency of spastic finger flexors) or on increasing kinesthetic awareness & recruitment of motor units in weak hypoactive muscles (e.g. wrist/ forearm extensor muscles).Reported benefits include improvements in range of motion, voluntary control & function²⁰⁻²⁴.

Methods

A total of forty five stroke participants aged between 40 to 65 years were screened for the study out of which thirty participants were eligible based upon the inclusion and exclusion criteria and those agreed to participate in the study. Two of these participants were dropped out of the study as they lost follow-up. Study group had 14 participants where as Control group had 14 participants. Criteria's for inclusion in this study were participants who had first ever M.C.A stroke, participants selected after two weeks of stroke and who had voluntary control grade 1-3 according to Hand Voluntary Control Scale. Participants were excluded who had any other musculoskeletal, cardiac vascular related conditions, cardiac pacemaker, impaired cognitive function (Mini Mental Scale), sensory loss over the hemiplegic hand.

Outcome measures

The outcome measures used in this study were Action research Arm Test and Voluntary control Grading Scale. It assesses a participant's ability to handle objects differing in size, weight and shape. The ARAT consists of 19 items grouped into four subscales: grasp, grip, pinch, and gross movement. Each subscale constitutes a hierarchical Guttman scale, which means that all items are ordered according to ascending difficulty. In the ARAT, if the participants succeed in completing the most difficult item in a subscale thus suggest he/she will succeeded in the easier items for that same subscale. Similarly, failure on an item suggests the participant will be unable to complete the remaining more challenging items in the subscale.²⁵

The voluntary control of hand was measured with the help of the voluntary control grading scale.

Procedure

The study received approval from Ethical Committee of Pravara Institute of Medical Sciences Deemed University, Loni 413736, Maharashtra, India. Participants were screened and based on the selection criteria they were included in the study. Those willing to participate were briefed about the nature of the study and effect of the intervention in the language best understood by them and the treatment was demonstrated to them and Written Informed Consent was obtained. All participants were given pre training for EMG biofeedback that how it feels after electrode placement. Explained the all procedure to the participants then randomly selected the participants for both Group A & Group B group. Study group received EMG Biofeedback along with conventional physiotherapy management. One electrode placed over the Extensor

Digitorum communis and Extensor Carpi muscle belly & another one place over the origin of the muscle. Asked the participants to do the wrist extension by giving visual feedback and at the same time asked to achieve the goal or asked to take the points which shown over the screen & it was for 20 min. Total treatment duration 40 minutes, 1 session / day, 5 session / week, total duration 4 weeks. Control group received squeeze ball exercise, putty exercise, cable wrist curl exercise, wrist circles exercise, finger ladder, cable roller exercise for 40 min / day. 5 days in a week for 4 weeks.




Figure 1 Flow diagram showing the procedure used in the study



Fig.2



Fig.3

Results

Statistical analysis was done by Graph Pad In Stat software using various statistical measures such as a mean, standard deviation (SD) and tests of significance such as unpaired 't' test. The results were concluded to be statistically significant with $p < 0.05$ and highly significant with $p < 0.01$. Unpaired 't' test was used to compare differences between the two groups i.e. the study group and the control group. No significant difference in baseline characteristics (Table 1). The results of this study were analyzed in terms of hand function and voluntary control indicated by improvement in ARA test and Voluntary control grading scale and comparison was made

between the first, 2nd week and after 4th week readings. The study results showed that EMG Biofeedback along with conventional physiotherapy are more effective than only conventional physiotherapy on improving hand function in hemiplegic participants as measured by Action Research Arm test and Voluntary control grading scale.

Table 1: Demographic profile and clinical data of participants of both the groups

	Group A (Study Group)	Group B (Control Group)	'p' value
Age (years)	52.93±5.91	55.26±4.23	0.22
Gender (M/F)	12(80%) /3(20%)	10(66.6%) /5(33.3%)	-
Duration after stroke (days)	2.466±0.516	2.866±0.833	0.125
Side of hemiplegia Rt/Lt	8 (53.3%) / 7(46.6%)	9(60%) / 6(40%)	-

Table: 2 Shows difference between pre and post mean values of Action Research Arm test of group A and group B.

Action Research Arm Test	Group A	Group B	't' value	'p' value	Results
	Mean ± SD	Mean ± SD			
After 2 nd week	2.357±1.277	1.285±2.268	1.540	0.135	Not significant
After 4 th week	5.857±3.159	1.714±2.268	3.986	0.05	Extremely significant




Fig 4 Graph showing pre and post values of Action Research Arm Test of group A and group B.

Table: 3 shows difference between Pre and post mean values of voluntary control grading scale of both group A and group B

Voluntary control grading scale	Group A	Group B	't' value	'p' value	Results
	Mean ± SD	Mean ± SD			
After 2 nd week	0.571±0.513	0.571±0.513	0.000	0.999	Not significant
After 4 th week	2±0.679	1±0.554	4.266	0.02	Extremely significant




Fig.4 Graph showing pre and post values of Voluntary Control Grading Scale of group A and group B.

Discussion

Study group showed greater improvement in improving hand function and voluntary control of hand in hemiplegic stroke participants. Improvement showed because of EMG Biofeedback and conventional physiotherapy which alter motor unit activity based on augmented audio & visual feedback information. Training can focus on voluntary inhibition of spastic muscles (e.g. reducing firing frequency of spastic finger flexors) or on increasing kinesthetic awareness & recruitment of motor units in weak hypoactive muscles (e.g. wrist/ forearm extensor muscles).²⁰⁻²⁴

Motor recovery in the arm was found to be significantly better in the study group. Therefore this study concluded that the effectiveness of the Biofeedback therapy could be attributed mainly to the repetitive stimulation of muscle activity in the arm. These results agree with those from two recent studies,^{27,28} in which active repetitive motor training of hand and fingers have proven to be effective. There is growing evidence that muscle weakness rather than spasticity plays a dominant role in impairment of active voluntary movements.^{29,30} Similar results were found in a favor of EMG Biofeedback in a study done by Armagan O et al (2003), conducted a study on “Electromyographic Biofeedback in the treatment of the Hemiplegic Hand” with 27 patients & concluded that the potential benefits of EMG biofeedback in conjunction with neurophysiologic rehabilitation technique to maximize the hand function in hemiplegic patients. Thus the clinical implication is the EMG Biofeedback along with conventional physiotherapy has significant effect on hand function in hemiplegic stroke participants when compared to control group. After 4th weeks of EMG Biofeedback session improvement was noted but at the 2nd weeks there was no significant effect of EMG Biofeedback along with conventional physiotherapy on hand function in hemiplegic stroke participants. A evidence from studies suggests that using EMG Biofeedback in combination with conventional physiotherapy may result in improvements in motor power beyond those of conventional physiotherapy alone. The limitations of this present study focused on only in MCA stroke participants as well as only subacute stage hemiplegic participants and ARA test and voluntary control grading scale both outcome measures were depends on the condition, the severity of disability of participants.

Conclusion

On the basis of this study it can be concluded that EMG Biofeedback along with conventional physiotherapy can be used on improving hand function and voluntary control of hand in hemiplegic stroke participants.

Acknowledgements

Ethical approval: Ethical Committee of Pravara Institute of Medical Sciences, Loni, Maharashtra state, India. (PMT/PIMS/RC/2011/08)

Funding: No funding.

References

1. Sims NR, Muyderman H (September 2009). "Mitochondria, oxidative metabolism and cell death in stroke". *Biochimica et Biophysica Acta* 1802 (1): 80–91. doi:10.1016/j.bbadis.2009.09.003. PMID 19751827.
2. Donnan GA, Fisher M, Macleod M, Davis SM (May 2008). "Stroke". *Lancet* 371 (9624): 1612–23. doi:10.1016/S0140-6736(08)60694-7. PMID 18468545.
3. World Health Organisation (1978). *Cerebrovascular Disorders (Offset Publications)*. Geneva: World Health Organization. ISBN 9241700432. OCLC 4757533.
4. American Heart Association (2005), Heart and Stroke Statistical -2005 Update. American Heart Association, Dallas
5. Post Stroke Rehabilitation Guideline Panel : Post Stroke Rehabilitation Clinical Practice Guideline. May 1995.
6. Brain stroke third biggest killer in India, health.indiatimes.com/articleshow /1148565.cms.
7. Barnett H, Mohr JP, Stein B, et al. (1992), *Stroke: Pathophysiology, Diagnosis and Management. 2nd ed.* London, England: Churchill Livingstone; 360-405.
8. Divyanshu Dubey, Anshudha Sawhney, Arvind Kavishwar, Sonjjay Pande, Devashish Dubey, (2011), A study of anatomical, seasonal and diurnal variation in the occurrence of Ischemic Stroke. *International Journal of Collaborative Research on Internal Medicine & public Health*. Vol. 3 No. 10
9. Schmidley JW, Messing RO. (1984), Agitated confusional states in patients with right hemisphere infarctions. *Stroke*. Sep-Oct 15(5):883-5.
10. Mohr JP, Rubinstein LV, Kase CS, et al. (1984), Hemiparesis profiles in stroke: the NINCDS stroke data bank. *Presented at the Annual Meeting of the American Neurological Association*. Baltimore, Md:
11. The Internet Stroke Center of Washington University: Blood Vessels of the Brain
12. Victor and Ropper, (2001) *Adam's and Victor's Principles of Neurology*
13. Antoni Fauci (ed); (2008) *Harrison's Principles of Internal Medicine*, 17th Edition.
14. Luci Fuscaldi Teixeira-Salmela, Sandra Jean Olney, Sylvie Nadeau, J. Brenda Brouwer, (1999), Muscle Strengthening and Physical Conditioning to Reduce Impairment and Disability in Chronic Stroke Survivors. *Arch Phys Med Rehabil* 80:1211-8.
15. Bobath, B. (1969), The treatment of neuromuscular disorders by improving patterns of coordination. *Physiotherapy*, 55:1,
16. Carvalho LMG. Terapia ocupacional na reabilitação de pacientes neurológicos adultos. In: DeCarlo M, Luzo M. (2004), *Terapia Ocupacional – Reabilitação Física e Contextos Hospitalares*. São Paulo: Editora Roca.
17. Dobkin BH. (2004), Strategies for stroke rehabilitation. *Lancet Neurol*. 3:528–536.
18. Gilroy, J., (2000), *Basic Neurology*, ed 3, McGraw-Hill, New York,
19. Wolf, S, and Binder-Macieod, S., (1983), Electromyographic biofeedback applications to the hemiplegic patients- changes in lower extremity neuromuscular and functional status . *Phys Ther* 63:1404,
20. Prevo, A, et al. (1982), Effect of EMG feedback on paretic muscles and abnormal cocontraction in the hemiplegic arm, compared with conventional physical therapy. *Scand J Rehabil Med* 14:121,
21. Tries, J. (1989), EMG Biofeedback for the treatment of upper extremity dysfunction: Can it be effective? *Biofeedback self Regul* 14(1):21.
22. Schleenbaker, RE and Mainous, AG. (1993), Electromyographic biofeedback for neuromuscular education in the hemiplegic stroke patient: A meta-analysis. *Arch Phys Med Rehabil* 74(12): 1301.
23. Glantz, M, et al (1995), Biofeedback therapy in post stroke rehabilitation: A meta-analysis of the randomised controlled trials. *Arch Phys Ther* 76:508
24. Brudny J, Korein J, Grynbaum BB, Belandres PV, Gianutsos JG. (1979), Helping hemiparetics to help themselves. Sensory feedback therapy. *JAMA*. 241:814–818.

25. Greenberg S, Fowler RS Jr. (1980), Kinesthetic biofeedback: a treatment modality for elbow range of motion in hemiplegia. *Am J Occup Ther.* 34:738–743.
26. Moreland J, Ann Thomson MA. (1994), Efficacy of Electromyographic Biofeedback Compared With Conventional Physical Therapy for Upper-Extremity Function in Patients Following Stroke: A Research Overview and Meta-analysis. *Phys Ther.* 74(6): 534-543
27. Bütetisch C, Hummelsheim H, Denzler P, Mauritz KH. (1995), Repetitive training of isolated movements improves the outcome of motor rehabilitation of the centrally paretic hand. *J Neurol Sci.* 130:59–68
28. Fellows SJ, Gandevia SC, Spina PJ. (1994), Voluntary movement at the elbow in spastic hemiparesis. *Ann Neurol.* 36:397–407
29. Wolf SL, LeCraw DE, Barton LA, Jann BB. (1989), Forced use of hemiplegic upper extremities to reverse the effect of learned nonuse among chronic stroke and head-injured patients. *Exp Neurol.* 104:125–132
30. Taub E, Miller NE, Novack TA, Cook EW, Fleming WC, Nepomuceno CS, Connell JS, Crago JE. (1993), Technique to improve chronic motor deficit after stroke. *Arch Phys Med Rehabil.* 74:347–354.